

Publications List
1976~2020

239. Iwasaki Y, Ikemura T, Kurokawa k, Okada N.
Implication of a new function of human tDNAs in chromatin organization.
Scientific Reports 10, 17440 (2020)
238. Vu TD, Iwasaki Y, Oshima K, Nikaido M, Chiu MT, Okada N.
A unique neurogenomic state emerges after aggressive confrontations in males of the fish *Betta splendens*.
bioRxiv 10,237586 (2020)
237. Vu TD, Iwasaki Y, Shigenobu S, Maruko A, Oshima K, Iioka E, Huang CL, Abe T, Tamaki S, Lin YW, Chen CK, Lu MY, Hojo M, Wang HV, Tzeng SF, Huang HJ, Kanai A, Gojobori T, Chiang TY, Sun HS, Li WH, Okada N.
Behavioral and brain- transcriptomic synchronization between the two opponents of a fighting pair of the fish *Betta splendens*.
PLOS Genetics 16(6), June (2020)
236. Okada N, Oshima K, Iwasaki Y, Maruko A, Iioka E, Vu T, Fujitsuka N, Nishi A, Sugiyama A, Nishiyama M, Kaneko A, Yamamoto M, Nishimura S.
Intron retention as a new marker of the pre-disease state and its recovery to the normal state by a traditional Japanese multi-herbal medicine.
bioRxiv 2020.02.10.941435 (2020)
235. Takuno S, Miyagi R, Onami J, Takahashi-kariyazono S, Sato A, Tichy H, Nikaido M, Aibara M, Mizoiri S, Mrosso D.J.H, Mzighani S.I, Okada N, Terai Y.
Patterns of genomic differentiation between two Lake Victoria cichlid species, *Haplochromis pyrrhocephalus* and *H. sp. 'macula'*.
BMC Evolutionary Biology 19(68), March (2019)
234. Sagai T, Amano T, Maeno A, Kimura T, Nakamoto M, Takehana Y, Naruse K, Okada N, Kiyonari H, Shiroishi T.
Evolution of Shh endoderm enhancers during morphological transition from ventral lungs to dorsal gas bladder.
Nature Communications 8 (14300), Feb (2017)
233. Terai Y, Miyagi R, Aibara M, Mizoiri S, Imai H, Okitsu T, Wada A, Takahashi-Kariyazono S, Sato A, Herbert T, Hillary D.J., Semvua I.M, Okada N.
Visual adaptation in Lake Victoria cichlid fishes: depth-related

variation of color and scotopic opsins in species from sand/mud bottoms.

BMC Evolutionary Biology 17, Article number:200 (2017)

232. Hori M, Nakajima M, Hata H, Yasugi M, Takahashi S, Nakae M, Yamaoka K, Kohada M, Kitamura J, Maehata M, Tanaka H, Okada N, Takeuchi Y.
Laterality is Universal Among Fishes but Increasingly Cryptic Among Derived Groups.
Zoological Science 2017 Aug ; 34(4):267-274 (2017)
231. Otsu M, Kajikawa M, Okada N, Kawai G.
Solution structure of a reverse transcriptase recognition site of a LINE RNA from zebrafish.
The Journal of Biochemistry, Volume 162, Issue 4, October 2017, Pages 279-285 (2017)
230. Nishihara H (12 coauthors) and Okada N.
Coordinately Co-opted Multiple Transposable Elements Constitute an Enhancer for *wnt5a* Expression in the Mammalian Secondary Palate.
PLOS Genetics journal pgen.1006380. (2016)
229. Nishihara H, Plazzi F, Passamonti M, Okada N.
MetaSINEs: Broad Distribution of a Novel SINE Superfamily in Animals.
Genome Biology and Evolution, Volume 8, Issue 3, Pages 528-539 (2016)
228. Tatsumi N, Kobayashi R, Yano T, Noda M, Fujimura K, Okada N, Okabe M.
Molecular developmental mechanism in polypterid fish provides insight into the origin of vertebrate lungs.
Scientific Reports 6, Article Number: 30580 (2016)
227. Kuramoto T, Nishihara H, Watanabe M, Okada N.
Determining the Position of Storks on the Phylogenetic Tree of Waterbirds by Retroposon Insertion Analysis.
Genome Biology and Evolution. Pages 3180–3189(2015)
226. Yamaguchi K, Kajikawa M, Okada N.
LINE retrotransposition and host DNA repair machinery.
Mobile Genetic Elements Volume 5 2015, Issue 6 Pages 92-97. Nov.2015 (2015)
225. Kudo Y, Nikaido M, Kondo A, Suzuki H, Yoshida K, Kikuchi K,

Okada N.

A microsatellite-based genetic linkage map and putative sex-determining genomic regions in Lake Victoria cichlids.

Gene Volume 560, Issue 2, Pages 156-164 (2015)

224. Suzuki H, Nikaido M, Hagino-Yamagishi K, Okada N.
Distinct functions of two olfactory marker protein genes derived from teleost-specific whole genome duplication.
BMC Evolutionary Biology 15, Article number:245 (2015)
223. Yamaguchi K, Kajikawa M, Okada N.
Integrated mechanism for the generation of the 5' junctions of LINE inserts
Nucleic Acids Research 42(21) 13269 -13279 (2014)
222. Shirai K, Inomata N, Mizoiri S, Aibara M, Terai Y, Okada N, Tachida H.
High prevalence of non-synonymous substitutions in mtDNA of cichlid fishes from Lake Victoria.
Gene Volume 552, Issue 2, 1 December 2014, Pages 239-245 (2014)
221. Brawand D, (45 coauthors), Nikaido M, Nishihara H, (18 coauthors), Okada N, (10coauthors) and Palma F.
The genomic substrate for adaptive radiation in African cichlid fish.
Nature (article) 513:375-381. (2014)
220. Hayashi Y, Kajikawa M, Matsumoto T, Okada N.
Mechanism by which a LINE protein recognizes its 3 tail RNA.
Nucleic Acids Research, Volume 42, Issue 16, 15 September 2014, Pages 10605–10617(2014)
219. Nikaido M, Ota T, Hirata T, Suzuki H, Satta Y, Aibara M, Mzighani SI, Sturmbauer C, Hagino-Yamagishi K, Okada N.
Multiple episodic evolution events in V1R receptor genes of East-African cichlids.
Genome Biology and Evolution Advance Access published May 6, 2014
doi:10.1093/gbe/evu086 (2014)
218. Matsumoto T, Terai Y, Okada N, Tachida.
Sensory drive speciation and patterns of variation at selectively neutral genes.
Evolutionary Ecology 28, 591-609 (2014)
217. Iwasaki Y, Abe T, Okada N, Wada K, Wada Y, Ikemura T.
Evolutionary Changes in Vertebrate Genome Signatures with Special

Focus on Coelacanth.

DNA Research, Volume 21, Issue 5, October 2014,

Pages 459-467(2014)

216. Kuroiwa A, Terai Y, Kobayashi N, Yoshida K, Suzuki M, Nakanishi A, Matsuda Y, Watanabe M, Okada N.
Construction of Chromosome Markers from the Lake Victoria Cichlid *Paralabidochromis chilotes* and Their Application to Comparative Mapping.
Cytogenetic and Genome Research 2014; 142:112-120 (2014)
215. Kawashima Y, Nishihara H, Akasaki T, Nikaido M, Tsuchiya K, Sagawa S, Okada N.
The complete mitochondrial genomes of deep-sea squid (*Bathyteuthis abyssicola*), bob-tail squid (*Semirossia patagonica*) and four giant cuttlefish (*Sepia apama*, *S. latimanus*, *S. lycidas* and *S. pharaonis*), and their application to the phylogenetic analysis of Decapodiformes.
Molecular Phylogenetics and Evolution Volume 69, Issue 3, December 2013, Pages 980-993(2013)
214. Takeda M, Kusumi J, Mizoiri S, Aibara M, Mzighani SI, Sato T, Terai Y, Okada N, Tachida H.
Genetic Structure of Pelagic and Littoral Cichlid Fishes from Lake Victoria.
PLOS ONE | Volume 8 | Issue 9 | e74088 (2013)
213. Terasaki N, Goodier JL, Cheung LE, Wang YJ, Kajikawa M, Kazazian Jr. H, Okada N.
In Vitro Screening for Compounds That Enhance Human L1 Mobilization.
PLOS ONE | Volume 8 | Issue 9 | e74629 (2013)
212. Nikaido M, Suzuki H, Toyoda A, Fujiyama A, Hagino-Yamagishi K, Kocher TD, Carleton K, Okada N.
Lineage-Specific Expansion of Vomeronasal Type 2 Receptor-like (OlfC) Genes in Cichlids May Contribute to Diversification of Amino Acid Detection Systems.
Genome Biology and Evolution Advance Access published March 15, 2013 doi:10.1093/gbe/evt041(2013)
211. Nikaido M (27 coauthors) Okada,N.
Coelacanth genomes reveal signatures for evolutionary transition from water to land.
Genome Res. 1, October (2013)
210. Yoshioka Y, Takahashi Y, Matsumoto S, Kojima S, Matsuoka K, Nakamura K, Ohoshima K, Okada N, Machida Y.

Mechanisms of T-DNA transfer and integration into plant chromosomes : role of *vir B*, *vir D4* and *vir E2* and a short interspersed repetitive element (SINE) from tobacco.
Molecular Mechanisms of Bacterial Virulence(2012)

209. Kajikawa M, Yamaguchi K, Okada N.
A new mechanism to ensure integration during LINE retrotransposition: A suggestion from analyses of the 5' extra nucleotides.
Gene. Mar 1 Epub ahead of print (2012)
208. Kajikawa M, Sugano T, Sakurai R, Okada N.
Low dependency of retrotransposition on the ORF1 protein of the zebrafish LINE, ZfL2-1.
Gene. May 10;499(1):41-7 (2012)
207. Ota T, Nikaido M, Suzuki H, Hagino-Yamaguchi K, Okada N.
Characterization of V1R receptor (*ora*) genes in Lake Victoria cichlids.
Gene May 15;499(2):273-9 (2012)
206. Nakamura M, Kajikawa M, Okada N.
Self-interaction, nucleic acid binding, and nucleic acid chaperone activities are unexpectedly retained in the unique ORF1p of zebrafish. LINE. Mol. Cell Biol. Jan 32 (2): 458-69 (2012)
205. Nakanishi A, Kobayashi N, Suzuki-Hirano A, Nishihara H, Sasaki T, Hirakawa M, Sumiyama K, Shimogori T, Okada N.
A SINE-derived element constitutes a unique modular enhancer for mammalian diencephalic *Fgf8*.
PLOS One 2012; 7(8)e43785 (2012)
204. Higasa K, Nikaido M, Saito TL, Yoshimura J, Suzuki Y, Suzuki H, Nishihara H, Aibara M, Ngatunga BP, Kalombo HW, Sugano S, Morishita S, Okada N.
Extremely slow rate of evolution in the HOX cluster revealed by comparison between Tanzanian and Indonesian coelacanths.
Gene. Jun 12. Epub ahead of print (2012)
203. Miyagi R, Terai Y, Aibara M, Sugawara T, Imai H, Tachida H, Mzighani SI, Okitsu T, Wada A, Okada N.
Correlation between nuptial colors and visual sensitivities tuned by opsins leads to species richness in sympatric Lake Victoria cichlid fishes.
Molecular Biology and Evolution, Volume 29, Issue 11, November 2012, Pages 3281-3296(2012)

202. Yuzawa Y, Nishihara H, Haraguchi T, Masuda S, Shimojima M, Shimoyama A, Yuasa H, Okada N, Ohta H.
Phylogeny of Galactolipid Synthase Homologs Together with their Enzymatic Analyses Revealed a Possible Origin and Divergence Time for Photosynthetic Membrane Biogenesis.
Author Notes DNA Research. Volume 19, Issue 1, February (2012)
201. Nikaido M, Sasaki T, Emerson J.J, Aibara M, Mzighani S.I, Budeba Y.L, Ngatunga B.P, Iwata M, Abe Y, Li W.H, Okada N.
Genetically distinct coelacanth population off the northern Tanzanian coast.
Proceedings of the National Academy of Sciences of the United States of America. 108 (44) 18009-18013 Nov.2011(2011)
200. Yoshida K, Terai Y, Mizoiri S, Aibara M, Nishihara H, Watanabe M, Kuroiwa A, Hirai H, Hirai Y, Matsuda Y, Okada N.
B chromosomes have a functional effect on female sex determination in lake victoria cichlid fishes.
PLOS Genet. 2011 ug;7(8): e1002203. (2011)
199. Tashiro K, Teissier A, Kobayashi N, Nakanishi A, Sasaki T, Yan K, Tarabykin V, Vigier L, Sumiyama K, Hirakawa M, Nishihara H, Pierani A, Okada N.
A mammalian conserved element derived from SINE displays enhancer properties recapitulating Satb2 expression in early-born callosal projection neurons.
PLOS One. 6 e28497 (2011)
198. Nagai H, Terai Y, Sugawara T, Imai H, Nishihara H, Hori M, Okada N.
Reverse evolution in RH1 for adaptation of cichlids to water depth in Lake Tanganyika.
Mol Biol Evol. Dec 20. (2010) [Epub ahead of print]
197. Okada N, Sasaki T, Shimogori T, Nishihara H.
Emergence of mammals by emergency: exaptation.
Genes Cells. Aug;15(8):801-12. Epub Jul 14. (2010)
196. Murata Y, Tamura M, Aita Y, Fujimura K, Murakami Y, Okabe M, Okada N, Tanaka M.
Allometric growth of the trunk leads to the rostral shift of the pelvic fin in teleost fishes.
Dev Biol. Nov 1;347(1):236-45. Epub Aug 6. (2010)
195. Akasaki T, Nikaido M, Nishihara H, Tsuchiya K, Segawa S, Okada N.
Characterization of a novel SINE superfamily from invertebrates:

"Ceph-SINEs" from the genomes of squids and cuttlefish.
Gene. Apr 15;454(1-2):8-19. (2010)

194. Sugawara T, Imai H, Nikaido M, Imamoto Y, Okada N
Vertebrate Rhodopsin Adaptation to Dim Light via Rapid Meta-II
Intermediate Formation. *Mol Biol Evol*. Mar;27(3):506-19. (2010)
193. Mzighani SI, Nikaido M, Takeda M, Seehausen O, Budeba YL,
Ngatunga BP, Katunzi EFB, Aibara M, Mizoiri S, Sato T, Tachida H,
Okada N.
Genetic Variation and Demographic History of the *Laparogramma*
Cichlid Species Group of Lake Victoria-An Analysis Based on SINEs
and Mitochondrial DNA. *Gene*. Jan 15;450(1-2):39-47. (2010)
192. Suzuki J, Yamaguchi K, Kajikawa M, Ichiyanagi K, Adachi N, Koyama
H, Takeda S, Okada N.
Genetic evidence that the non-homologous end-joining repair pathway
is involved in LINE retrotransposition.
PLoS Genet. Apr;5(4):e1000461.(2009)
191. Ohara H, Nikaido M, Date-Ito A, Mogi K, Okamura H, Okada N,
Takeuchi Y, Mori Y, Hagino-Yamagishi K.
Conserved repertoire of orthologous vomeronasal type 1 receptor genes
in ruminant species.
BMC Evol. Biol. 9(1):233. (2009)
190. Okada N.
Special issue on phylogenomics and its future
Gene Jul 15;441(1-2):1-2 (2009) [Editorial]
189. Kojima K.K, and Okada N.
mRNA retrotransposition coupled with 5' inversion as a possible source
of new genes.
Mol Biol Evol. 26 (6): 1405-20 (2009)
188. Nishihara H, Maruyama S, Okada N.
Retroposon analysis and recent geological data suggest near-
simultaneous divergence of the three superorders of mammals.
Proc Natl Acad Sci USA. Mar 31;106(13):5235-40 (2009) [Abstract]
[OPEN ACCESS]
187. Kobayashi N, Watanabe M, Horiike T, Kohara Y, Okada N.
Extensive analysis of EST sequences reveals that all cichlid species in
Lake Victoria share almost identical transcript sets.
Gene. Jul 15;441(1-2):187-91 (2009)

186. Hirakawa M, Nishihara H, Kanehisa M, Okada N.
Characterization and evolutionary landscape of AmnSINE1 in Amniota genomes.
Gene Jul 15;441(1-2):100-10 (2009)
185. Maeda K, Takeda M, Kamiya K, Aibara M, Mzighani S.I, Nishida M, Mizoiri S, Sato T, Terai Y, Okada N, Tachida H.
Population structure of two closely related pelagic cichlids in Lake Victoria, *Haplochromis pyrrhocephalus* and *H. laparogramma*.
Gene Jul 15;441(1-2):67-73 (2009)
184. Murata Y, Yonezawa T, Kihara I, Kashiwamura T, Sugihara Y, Nikaido M, Okada N, Endo H, Hasegawa M.
Chronology of the extant African elephant species and case study of the species identification of the small African elephant with the molecular phylogenetic method.
Gene Jul 15;441(1-2):176-86 (2009)
183. Piskurek O, Nishihara H, Okada N.
The evolution of two partner LINE/SINE families and a full-length chromodomain-containing Ty3/Gypsy LTR element in the first reptilian genome of *Anolis carolinensis*.
Gene Jul 15;441(1-2):111-8 (2009)
182. Matveev V, Okada N.
Retroposons of salmonoid fishes (Actinopterygii: Salmonoidei) and their evolution.
Gene. 434(1-2):16-28 (2009)
181. Seehausen O, Terai Y, Magalhaes I. S, Carleton K. L., Mrosso H. D. J, Miyagi R, van der Sluijs I, Schneider M. V, Maan M, Tachida H, Imai H, Okada N.
Speciation through sensory drive in cichlid fish.
Nature. Oct 2;455(7213) 620-6 (2008)
180. Fujimura K, Okada N
Shaping of the lower jaw bone during growth of Nile tilapia *Oreochromis niloticus* and a Lake Victoria cichlid *Haplochromis chilotes*: A geometric morphometric approach.
Develop. Growth Differ. Oct;50(8):653-63. (2008)
179. Nishihara H, Okada N.
Retroposons: Genetic Footprints on the Evolutionary Paths of Life.
Phylogenomics (Methods in Molecular Biology, vol. 422)
Chapter 13, Pages 201-225, Ed. Murphy, W. J., Humana Press. (2008)

178. Matsuura M, Nishihara H, Onimaru K, Kokubo N, Kuraku S, Kusakabe R, Okada N, Kuratani S, Tanaka M.
Identification of Four Engrailed Genes in the Japanese Lamprey, *Lethenteron japonicum*.
Dev. Dyn. Jun;237(6):1581-9. (2008)
177. Ichiyanagi K, Okada N.
Mobility Pathways for Vertebrate L1-, L2-, CR1-, and RTE-Clade Retrotransposons.
Mol Biol Evol. Jun;25(6):1148-57 (2008)
176. Fujimura K, Okada N
Bone development in the jaw of Nile tilapia *Oreochromis niloticus* (Pisces: Cichlidae)
Develop. Growth Differ. 50(5): 339-55 (2008)
175. Fujimura K, Terai Y, Ishiguro N, Miya M, Nishida M, Okada N
Heterotopy in the N-terminal region of Growth/Differentiation Factor 5 (GDF5) mature protein during teleost evolution
Mol Biol Evol 25(5): 797-800 (2008)
174. Sasaki T, Nishihara H, Hirakawa M, Fujimura K, Tanaka M, Kokubo N, Kimura-Yoshida C, Matsuo I, Sumiyama K, Saitou N, Shimogori T, Okada N.
Possible involvement of SINEs in mammalian-specific brain formation.
Proc Natl Acad Sci USA. v105(11), 4220-5 (2008) [OPEN ACCESS]
173. Maeda K, Takeshima H, Mizoiri S, Okada N, Nishida M, Tachida H.
Isolation and characterization of microsatellite loci in the cichlid fish in Lake Victoria, *Haplochromis chilotes*.
Molecular Ecology Resources 8(2) 428 - 430 (2008)
172. Munemasa M, Nikaido M, Nishihara H, Donnellan S, Austin CC, Okada N.
Newly discovered young CORE-SINEs in marsupial genomes.
Gene. Jan 15;407(1-2):176-85. (2008)
171. Nishihara H, Okada N, Hasegawa M.
Rooting the eutherian tree: the power and pitfalls of phylogenomics.
Genome Biol. Sep 21;8(9) R199 (2007)
170. Nishihara H, Kuno S, Nikaido M, Okada N.
MyrSINEs: a novel SINE family in the anteater genomes.
Gene. Oct 1;400(1-2):98-103. (2007)

169. Ichiyanagi K, Nishihara H, Duvernell D.D, Okada N.
Acquisition of Endonuclease Specificity During Evolution of L1 Retrotransposon.
Mol Biol Evol. Sep;24(9):2009-15 (2007)
168. Watanabe M, Hiraide K, Okada N.
Functional diversification of kir7.1 in cichlids accelerated by gene duplication.
Gene Sep;399(1):46-52 (2007)
167. Piskurek O, Okada N.
Poxviruses as possible vectors for horizontal transfer of retroposons from reptiles to mammals.
Proc Natl Acad Sci U S A. Jul 17;104(29):12046-51 (2007) [OPEN ACCESS]
166. Matveev V, Nishihara H, Okada N.
Novel SINE Families from Salmon Validate Parahucho (Salmonidae) as a Distinct Genus and Evidence that SINEs Can Incorporate LINE-related 3'-Tails of Other SINEs.
Mol Biol Evol. Aug;24(8):1656-66 (2007)
165. Onami J, Nikaido M, Mannen H, Okada N.
Genomic expansion of the Bov-A2 retroposon relating to phylogeny and breed management.
Mamm Genome. Mar;18(3):187-96. (2007)
164. Yonezawa T, Nikaido M, Kohno N, Fukumoto Y, Okada N, Hasegawa M.
Molecular phylogenetic study on the origin and evolution of Mustelidae.
Gene. Jul 1;396(1):1-12 (2007)
163. Xue B, Rooney AP, Kajikawa M, Okada N, Roelofs WL.
Novel sex pheromone desaturases in the genomes of corn borers generated through gene duplication and retroposon fusion.
Proc Natl Acad Sci U S A. Mar 13;104(11):4467-72 (2007) [OPEN ACCESS]
162. Fujimura K, Okada N.
Development of the embryo, larva and early juvenile of Nile tilapia *Oreochromis niloticus* (Pisces: Cichlidae).
Developmental staging system
Develop. Growth Differ. 49 (4), 301-324 (2007)

161. Nikaido M, Piskurek O, Okada N.
Toothed whale monophyly reassessed by SINE insertion analysis: The absence of lineage sorting effects suggests a small population of a common ancestral species.
Mol. Phylogenet. Evol. 43:216-224 (2007)
160. Honda H, Ichiyanagi K, Suzuki J, Ono T, Koyama H, Kajikawa M, Okada N.
A new system for analyzing LINE retrotransposition in the chicken DT40 cell line widely used for reverse genetics.
GENE, Jun 15;395(1-2):116-24 (2007)
159. Sasaki T, Sato T, Miura S, Bwathondi PO, Ngatunga BP, Okada N.
Mitogenomic analysis for coelacanths (*Latimeria chalumnae*) caught in Tanzania.
Gene. 389 73-79 (2007)
158. Babushok D.V, Ohshima K, Ostertag E.M, Chen X, Wang, Mandal P.K, Okada N, Abrams C.S, Kazazian Jr.H.H.
A novel testis ubiquitin-binding protein gene arose by exon shuffling in hominoids.
Genome Research 17(8) 1129 - 1138 (2007)
157. Ichiyanagi K, Nakajima R, Kajikawa M, Okada N.
Novel retrotransposon analysis reveals multiple mobility pathways dictated by hosts.
Genome Res. Jan;17(1):33-41 (2007)
156. Sasaki T, Yasukawa Y, Takahashi K, Miura S, Shedlock AM, Okada N.
Extensive morphological convergence and rapid radiation in the evolutionary history of the family Geoemydidae (old world pond turtles) revealed by SINE insertion analysis.
Syst Biol. 55(6),912-27 (2006)
155. Tamura M, Kajikawa M, Okada N.
Functional splice sites in a zebrafish LINE and their influence on zebrafish gene expression.
Gene. Apr 1;390(1-2):221-31 (2006)
154. Terai Y, Seehausen O, Sasaki T, Takahashi K, Mizoiri S, Sugawara T, Sato T, Watanabe M, Konijnendijk N, Mrosso HD, Tachida H, Imai H, Shichida Y, Okada N.
Divergent Selection on Opsins Drives Incipient Speciation in Lake Victoria Cichlids.
PLoS Biol. 4(12):e433 (2006)

153. Iwashita M, Watanabe M, Ishii M, Chen T, Johnson SL, Kurachi Y, Okada N, Kondo S.
Pigment Pattern in jaguar/obelix Zebrafish Is Caused by a Kir7.1 Mutation: Implications for the Regulation of Melanosome Movement.
PLoS Genet. Nov.24;2(11): e197 (2006)
152. Ichiyanagi K, Okada N.
Genomic alterations upon integration of zebrafish L1 elements revealed by the TANT method.
Gene. 15;383: 108-16. (2006)
151. Watanabe M, Iwashita M, Ishii M, Kurachi Y, Kawakami A, Kondo S, Okada N.
Spot pattern of leopard Danio is caused by mutation in the zebrafish connexin41.8 gene.
EMBO Rep. 7(9):893-7. (2006)
150. Nomura Y, Kajikawa M, Baba S, Nakazato S, Imai T, Sakamoto T, Okada N, Kawai G.
Solution structure and functional importance of a conserved RNA hairpin of eel LINE UnaL2.
Nucleic Acids Res. 34;18: 5184-93 (2006)
149. Sasaki T, Nikaido M, Wada S, Yamada T K, Cao Y, Hasegawa M, Okada N.
Balaenoptera omurai is a newly discovered baleen whale that represents an ancient evolutionary lineage.
Mol Phylogenet Evol. 41;40 :40-52 (2006)
148. Munemasa M, Nikaido M, Donnellan S, Austin C C, Okada N, Hasegawa M.
Phylogenetic analysis of diprotodontian marsupials based on complete mitochondrial genomes.
Genes Genet Syst. 81:181-91 (2006)
147. Piskurek O, Okada N.
Simple and complex SINEs: A brief critical comment.
Gene 375, 110 (2006)
146. Watanabe M, Nikaido M, Tsuda T T, Kobayashi T, Mindell D, Cao Y, Okada N, Hasegawa M.
New candidate species most closely related to penguins.
Gene 378, 65-73 (2006)

145. Nishihara H, Hasegawa M, Okada N.
Pegasoferae, an unexpected mammalian clade revealed by tracking ancient retroposon insertions.
Proc Natl Acad Sci USA. 103(26),9929-34 (2006) [OPEN ACCESS]
144. Nishihara H, Smit AF, Okada N.
Functional noncoding sequences derived from SINEs in the mammalian genome.
Genome Res. Jul;16(7):864-74 (2006)
143. Piskurek O, Austin CC, Okada N.
Sauria SINEs: Novel Short Interspersed Retroposable Elements That Are Widespread in Reptile Genomes.
J Mol Evol. 62(5), 630-44. (2006)
142. Nikaido M, Hamilton H, Makino H, Sasaki T, Takahashi K, Goto M, Kanda N, Pastene LA, Okada N.
Baleen Whale Phylogeny and a Past Extensive Radiation Event Revealed by SINE Insertion Analysis.
Mol Biol Evol. 23, :866-73 (2006)
141. Kobayashi N, Watanabe M, Kijimoto T, Fujimura K, Nakazawa M, Ikeo K, Kohara Y, Gojobori T, Okada N.
magp4 gene may contribute to the diversification of cichlid morphs and their speciation.
Gene 373, 126-33 (2006)
140. Akasaki T, Nikaido M, Tsuchiya K, Segawa S, Hasegawa M, Okada N.
Extensive mitochondrial gene arrangements in coleoid Cephalopoda and their phylogenetic implications.
Mol Phylogenet Evol. 38, 648-58 (2006)
139. Sugano T, Kajikawa M, Okada N.
Isolation and characterization of retrotransposition-competent LINEs from zebrafish.
Gene 365, 74-82. (2006)
138. Kawagoe-Takaki H, Nameki N, Kajikawa M, Okada N.
Probing the secondary structure of salmon SmaI SINE RNA.
Gene 365, 67-73. (2006)
137. Watanabe M, Nikaido M, Tsuda TT, Inoko H, Mindell DP, Murata K, Okada N.
The rise and fall of the CR1 subfamily in the lineage leading to

penguins.
Gene 365, 57-66. (2006)

136. Ohshima K, and Okada N.
SINEs and LINEs: symbionts of eukaryotic genomes with a common tail.
Cytogenet Genome Res. 110, 475-90. (2005)
135. Nishihara H, Satta Y, Nikaido M, Thewissen J G, Stanhope M J, and Okada N.
A Retroposon Analysis of Afrotherian Phylogeny.
Mol Biol Evol. 22(9), 1823-33. (2005)
134. Kijimoto T, Watanabe M, Fujimura K, Nakazawa M, Murakami Y, Kuratani S, Kohara Y, Gojobori T, and Okada N.
cimp1, A Novel Astacin Family Metalloproteinase Gene from East African Cichlids, Is Differentially Expressed Between Species During Growth.
Mol Biol Evol. 22(8), 1649-60. (2005)
133. Sugawara T, Terai Y, Imai H, Turner G F, Koblmuller S, Sturmbauer C, Shichida Y, and Okada N.
Parallelism of amino acid changes at the RH1 affecting spectral sensitivity among deep-water cichlids from Lakes Tanganyika and Malawi.
Proc. Natl. Acad. Sci. USA 102(15), 5448-53 (2005) [OPEN ACCESS]
132. Sasaki T, Nikaido M, Hamilton H, Goto M, Kato H, Kanda N, Pastene L A, Cao A, Fordyce R E, Hasegawa M, and Okada N.
Mitochondrial phylogenetics and evolution of mysticete whales.
Syst Biol. 54, 77-90 (2005)
131. Kajikawa M, Ichiyanagi K, Tanaka N, and Okada N.
Isolation and characterization of active LINE and SINEs from the eel.
Mol. Biol. Evol. 22, 673-82 (2005)
130. Segawa T, Miyamoto K, Ushida K, Agata K, Okada N, and Kohshima S.
Seasonal Change in Bacterial Flora and Biomass in Mountain Snow from the Tateyama Mountains, Japan, Analyzed by 16S rRNA Gene Sequencing and Real-Time PCR.
Applied and Environmental Microbiology 71(1), 123-30 (2005)
129. Sugie A, Terai Y, Ota R, and Okada N.
The evolution of genes for pigmentation in African cichlid fishes.
GENE 343(2), 337-46 (2004)

128. Watanabe M, Kobayashi N, Shin-i T, Tateno Y, Kohara Y, and Okada N.
Extensive analysis of ORF sequences from two different cichlid species in Lake Victoria provides molecular evidence for a recent radiation event of the Victoria species flock.
GENE 343(2), 263-9 (2004)
127. Baba S, Kajikawa M, Okada N, and Kawa G.
Solution structure of an RNA stem-loop derived from the 3' conserved region of eel LINE UnaL2.
RNA 10(9), 1380-7 (2004)
126. Shedlock A M, Takahashi K, and Okada N.
SINEs of speciation: tracking lineages with retroposons.
Trends in Ecology & Evolution 19(10), 545-53 (2004)
125. Imanishi T, et al.
Integrative Annotation of 21,037 Human Genes Validated by Full-Length cDNA Clones.
Plos Biology 2(6), 856-75 (2004)
124. Okada N, Shedlock A M, and Nikaido M.
Retroposon mapping in molecular systematics.
Methods. Mol. Biol. 260, 189-226 (2004)
123. Sasaki T, Takahashi K, Nikaido M, Miura S, Yasukawa Y, and Okada N.
First Application of the SINE (Short Interspersed Repetitive Element) Method to Infer Phylogenetic Relationships in Reptiles: An Example from the Turtle Superfamily Testudinoidea.
Mol. Biol. Evol. 21, 705-15 (2004)
122. Terai Y, Takezaki N, Mayer W E, Tichy H, Takahata N, Klein J, and Okada N.
Phylogenetic relationships among East African haplochromine fishes as revealed by short interspersed elements (SINEs).
J. Mol. Evol. 58, 64-78 (2004)
121. Kawai K, Nikaido M, Harada M, Matsumura S, Lin L.K, Yi Wu, Hasegawa M, Okada N.
Erratum: The status of the Japanese and East Asian bats of the genus *Myotis* (Vespertilionidae) based on mitochondrial sequences (*Molecular Phylogenetics and Evolution* (2003) 28 (297-307))
Molecular Phylogenetics and Evolution. 30(1)259(2004)

120. Ohshima K, Hattori M, Yada T, Gojobori T, Sakaki Y, and Okada N.
Whole-genome screening indicates a possible burst of formation of processed pseudogenes and Alu repeats by particular L1 subfamilies in ancestral primates.
Genome Biol. 4(11)R74, Epub 2003 Oct 28. (2003)
119. Terai Y, Morikawa N, Kawakami K, and Okada N.
The complexity of alternative splicing of hagoromo mRNA is increased in an explosively speciated lineage in East African cichlids.
Proc. Natl. Acad. Sci. U S A. Oct.28,100(22), 12798-803 (2003) [OPEN ACCESS]
118. Terai Y, Takahashi K, Nishida M, Sato T, and Okada N.
Using SINEs to Probe Ancient Explosive Speciation: "Hidden" Radiation of African Cichlids?
Mol. Biol. Evol. 20, 924-30 (2003)
117. Piskurek O, Nikaido M, Boeadi Baba M, and Okada N.
Unique mammalian tRNA-derived repetitive elements in Dermopterans: The t-SINE family and its retrotransposition through multiple sources.
Mol. Biol. Evol. 20, 1659-68 (2003)
116. Murata Y, Nikaido M, Sasaki T, Cao Y, Fukumoto Y, Hasegawa M, and Okada N.
Afrotherian phylogeny as inferred from complete mitochondrial genomes.
Mol. Phylogenet. Evol. 28, 253-60 (2003)
115. Nikaido M, Cao Y, Harada M, Okada N, and Hasegawa M.
Mitochondrial phylogeny of hedgehogs and monophyly of Eulipotyphla.
Mol. Phylogenet. Evol. 28, 276-84 (2003)
114. Kawai K, Nikaido M, Harada M, Matsumura S, Lin L K, Wu Y., Hasegawa M and Okada N.
The status of the Japanese and East Asian bats of the genus *Myotis* (Vespertilionidae) based on mitochondrial sequences.
Mol. Phylogenet. Evol. 28, 297-307 (2003)
113. Nikaido M, Nishihara H, Fukumoto Y, and Okada N.
Ancient SINEs from African Endemic Mammals
Mol. Biol. Evol. 20, 522-7 (2003)

112. Watanabe M, Kobayashi N, Fujiyama A, and Okada N.
Construction of a BAC library for *Haplochromis chilotes*, a cichlid fish from Lake Victoria.
Genes. Genet. Syst. 78, 103-5 (2003)
111. Nikaido M, Cao Y, Okada N, and Hasegawa M.
The phylogenetic relationships of insectivores with special reference to the lesser hedgehog tenrec as inferred from the complete sequence of their mitochondrial genome.
Genes. Genet. Syst. 78, 107-12 (2003)
110. Ishitani R, Nureki O, Nameki N, Okada N, Nishimura S, and Yokoyama S.
Alternative Tertiary Structure of tRNA for Recognition by a Posttranscriptional Modification Enzyme.
Cell 113, 383-94 (2003)
109. D Huchon, Nikaido M, Okada N.
Use of SINE retroposon in phylogeny.
M S-MEDECINE SCIENCES 18(12) 1276-1281(2002)
108. Pupko T, Huchon D, Cao Y, Okada N, and Hasegawa M.
Combining multiple data sets in a likelihood analysis: which models are the best?
Mol Biol Evol. 19(12), 2294-307 (2002)
107. Terai Y, Mayer W E, Klein J, Tichy H, and Okada N.
The effect of selection on a long wavelength-sensitive (LWS) opsin gene of Lake Victoria cichlid fishes.
Proc. Natl. Acad. Sci. USA 99, 15501-6 (2002) [OPEN ACCESS]
106. Kajikawa M, and Okada N.
LINEs Mobilize SINEs in the Eel through a Shared 3' Sequence.
Cell 111, 433-44 (2002)
105. Nishihara H, Terai Y, and Okada N.
Characterization of Novel Alu- and tRNA-Related SINEs from the Tree Shrew and Evolutionary Implications of Their Origins.
Mol. Biol. Evol. 19, 1964-72 (2002)
104. Sugawara T, Terai Y, and Okada N.
Natural selection of the rhodopsin gene during the adaptive radiation of East african great lakes cichlid fishes.
Mol. Biol. Evol. 19, 1807-11 (2002)

103. Terai Y, Morikawa N, and Okada N.
The evolution of the pro-domain of bone morphogenetic protein 4 (bmp4) in an explosively speciated lineage of East african cichlid fishes.
Mol. Biol. Evol. 19(9), 1628-32 (2002)
102. Kawai K, Nikaido M, Harada M, Matsumura S, Lin L K, Wu Y, Hasegawa M, and Okada N.
Intra- and Interfamily Relationships of Vespertilionidae Inferred by Various Molecular Markers Including SINE Insertion Data.
J. Mol. Evol. 55(3), 284-301 (2002)
101. Takahashi K, and Okada N.
Mosaic structure and retropositional dynamics during evolution of subfamilies of short interspersed elements in african cichlids.
Mol. Biol. Evol. 19(8), 1303-12 (2002)
100. Terai Y, Morikawa N, Kawakami K, and Okada N.
Accelerated Evolution of the Surface Amino Acids in the WD-Repeat Domain Encoded by the hageromo Gene in an Explosively Speciated Lineage of East African Cichlid Fishes.
Mol. Biol. Evol. 19(4), 574-8 (2002)
99. Ishitani R, Nureki O, Fukai S, Kijimoto T, Nameki N, Watanabe M, Kondo H, Sekine M, Okada N, Nishimura S, and Yokoyama S.
Crystal structure of archaeosine tRNA-guanine transglycosylase.
J. Mol. Biol. 318(3), 665-77 (2002)
98. Ogiwara I, Miya M, Ohshima K, and Okada N.
V-SINEs: a new superfamily of vertebrate SINEs that are widespread in vertebrate genomes and retain a strongly conserved segment within each repetitive unit.
Genome Res. 12, 316-24 (2002)
97. Ishitani R, Nureki O, Kijimoto T, Watanabe M, Kondo H, Nameki N, Okada N, Nishimura S, and Yokoyama S.
Crystallization and preliminary X-ray analysis of the archaeosine tRNA-guanine transglycosylase from *Pyrococcus horikoshii*.
Acta. Crystallogr. 57, 1659-62 (2001)
96. Nikaido M, Matsuno F, Abe H, Shimamura M, Hamilton H, Matsubayashi H, and Okada N.
Evolution of CHR-2 SINEs in cetartiodactyl genomes: possible evidence for the monophyletic origin of toothed whales.
Mamm. Genome. 12, 909-15 (2001)

95. Takahashi K, Terai Y, Nishida M, and Okada N.
Phylogenetic relationships and ancient incomplete lineage sorting among cichlid fishes in Lake Tanganyika as revealed by analysis of the insertion of retroposons.
Mol. Biol. Evol. 18, 2057-66 (2001)
94. Nikaido M, Kawai K, Cao Y, Harada M, Tomita S, Okada N, and Hasegawa M.
Maximum likelihood analysis of the complete mitochondrial genomes of eutherians and a reevaluation of the phylogeny of bats and insectivores.
J. Mol. Evol. 53, 508-16 (2001)
93. Takahashi K, Nishida M, Yuma M, and Okada N.
Retroposition of the AFC family of SINEs (short interspersed repetitive elements) before and during the adaptive radiation of cichlid fishes in Lake Malawi and related inferences about phylogeny.
J. Mol. Evol. 53, 496-507 (2001)
92. Nikaido M, Matsuno F, Hamilton H, Brownell R L Jr., Cao Y, Ding W, Zuoyan Z, Shedlock A M, Fordyce R E, Hasegawa M, and Okada N.
Retroposon analysis of major cetacean lineages: the monophyly of toothed whales and the paraphyly of river dolphins.
Proc. Natl. Acad. Sci. USA. 98(13), 7384-9 (2001) [OPEN ACCESS]
91. Watanabe M, Nameki N, Matsuo-Takasaki M, Nishimura S, and Okada N.
tRNA recognition of tRNA-guanine transglycosylase from a hyperthermophilic archaeon, *Pyrococcus horikoshii*.
J. Biol. Chem. 276(4), 2387-94 (2001)
90. Cao Y, Fujiwara M, Nikaido M, Okada N, and Hasegawa M.
Interordinal relationships and timescale of eutherian evolution as inferred from mitochondrial genome data.
GENE 259, 149-58 (2000)
89. Nikaido M, and Okada N.
CetSINEs and AREs are not SINEs but are parts of cetartiodactyl L1.
Mamm. Genome. 11 (12) , 1123-6 (2000)
88. Nikaido M, Harada M, Cao Y, Hasegawa M, and Okada N.
Monophyletic origin of the order chiroptera and its phylogenetic position among mammalia, as inferred from the complete sequence of the mitochondrial DNA of a Japanese megabat, the Ryukyu flying fox (*Pteropus dasymallus*).
J. Mol. Evol. 51(4), 318-28 (2000)

87. Lum J K, Nikaido M, Shimamura M, Shimodaira H, Shedlock A M, Okada N, and Hasegawa M.
Consistency of SINE insertion topology and flanking sequence tree: quantifying relationships among cetartiodactyls.
Mol. Biol. Evol. 17 (10) , 1417-24 (2000)
86. Shedlock A M, and Okada N.
SINE Insertions : Powerful Tools for Molecular Systematics.
BioEssays. 22(2), 148-60 (2000)
85. Shedlock A M, Milinkovitch M C, and Okada N.
SINE evolution, missing data, and the origin of whales.
Syst. Bio. 49(4), 808-17 (2000)
84. Waddell P J, Okada N, and Hasegawa M.
Towards resolving the interordinal relationships of placental mammals.
Syst. Biol. 48, 1-5 (1999)
83. Nikaido M, Rooney A P, and Okada N
Phylogenetic relationships among cetartiodactyls based on insertions of short and long interspersed elements: Hippopotamuses are the closest extant relatives of whales.
Proc. Natl. Acad. Sci. USA. 96, 10261-6 (1999) [OPEN ACCESS]
82. Shimamura M, Abe H, Nikaido M, Ohshima K, and Okada N.
Genealogy of families of SINEs in cetaceans and artiodactyls: The presence of a huge superfamily of tRNA Glu-derived families of SINEs.
Mol. Bio. Evol. 16(8), 1046-60 (1999)
81. Ogiwara I, Miya M, Ohshima K, and Okada N.
Retropositional parasitism of SINEs on LINEs: Identification of SINEs and LINEs in elasmobranchs.
Mol. Bio. Evol. Sep.16(9), 1238-50 (1999)
80. Murata S, Takasaki N, Okazaki T, Kobayasi T, Numachi K, Chang K H, and Okada N.
Molecular evidence from short interspersed elements (SINEs) that *Oncorhynchus masou* (cherry salmon) is monophyletic.
Can. J. Fish. Aquat. Sci. 55(8): 1864-70 (1998)
79. Terai Y, Takahashi K, and Okada N.
SINE cousins: The 3'-end tails of the two oldest and distantly related families of SINEs are descended from the 3' ends of LINEs with the

- same genealogical origin.
Mol. Biol. Evol. 15, 1460-71 (1998)
78. Shimamura M, Nikaido M, Ohshima K, and Okada N.
A SINE that acquired a role in signal transduction during evolution.
Mol. Biol. Evol. 15, 923-5 (1998)
77. Hamada M, Takasaki N, Reist J D, DeCicco A L, Goto A, and Okada N.
Detection of the on-going sorting of ancestrally polymorphic SINEs
towards fixation or loss in populations of two species of charr during
speciation.
Genetics 150(1), 301-11 (1998)
76. Takahasi I, Nobukuni T, Ohmori H, Kobayashi M, Tanaka S, Ohshima
K, Okada N, Masui T, Hashimoto K, and Iwashita S.
Existence of a bovine LINE repetitive insert that appears in the cDNA of
bovine protein BCNT in ruminant, but not in human, genomes.
GENE 211, 387-94 (1998).
75. Takahashi K, Terai Y, Nishida M, and Okada N.
A novel family of short interspersed repetitive elements (SINEs) from
cichlids: the patterns of insertion of SINEs at orthologous loci support
the proposed monophyly of four major groups of cichlid fishes in Lake
Tanganyika.
Mol. Biol. Evol. 15(4), 391-407 (1998)
74. Hamada M, Himberg M, Bodalu R A, Reist J D, and Okada N.
Monophyletic origin of the genera *stenodus* and *coregonus* as inferred
from an analysis of the insertion of SINEs (short interspersed repetitive
elements).
Biology and Management of Coregonid Fishes. 50, 383-9 (1998)
73. Cao Y, Waddell P J, Okada N, and Hasegawa M.
The complete mitochondrial DNA sequence of the shark *Mustelus
manazo*: evaluating rooting contradictions to living bony vertebrates .
Mol. Biol. Evol. 15, 1637-46 (1998)
72. Cao Y, Janke A., Waddell P J., Westerman M, Takenaka O, Murata S,
Okada N, Paabo S, and Hasegawa M.
Conflict among individual mitochondrial proteins in resolving the
phylogeny of eutherian orders.
J. Mol. Evol. 47, 307-22 (1998)
71. Watanabe M, Matsuo M, Tanaka S, Akimoto H, Asahi S, Nishimura S,
Katze J R, Hashizume T, Crain P F, McCloskey J A, and Okada N.

Biosynthesis of archaeosine, a novel derivative of 7-deazaguanosine specific to archaeal tRNA, proceeds via a pathway involving base replacement on the tRNA polynucleotide chain.

J. Biol. Chem. 272, 20146-51 (1997)

70. Okada N, Hamada M, Ogiwara I, and Ohshima K.
SINEs and LINEs share common 3' sequences: a review.
GENE 205(1-2), 229-43 (1997)
69. Shimamura M, Yasue H, Ohshima K, Abe H, Kato H, Kishiro T, Goto M, Munechika I, and Okada N.
Molecular evidence that whales form a clade within even-toed ungulates.
Nature 388 (6643) , 666-70 (1997)
68. Ishida K, Cao Y, Hasegawa M, Okada N, and Hara Y.
The origin of chlorarachniophyte plastids, as inferred from phylogenetic comparisons of amino acid sequences of EF-Tu.
J. Mol. Evol. 45, 682-7 (1997)
67. Hamada M, Kido Y, Himberg M, Hasegawa M, and Okada N.
A newly isolated family of short interspersed repetitive elements (SINEs) in coregonid fishes (whitefishes) with sequences that are almost identical to those of the Sma I family of repeats: possible evidence for the horizontal transfer of SINEs.
Genetics 146(1), 355-67 (1997)
66. Takasaki N, Yamaki T, Hamada M, Park L, and Okada N.
The salmon Sma I family of SINEs: interspecific and intraspecific polymorphism of the insertion of SINEs in the genomes of chum and pink salmon.
Genetics 146(1), 369-80 (1997)
65. Kajikawa M, Ohshima K, and Okada N.
Determination of the entire sequence of turtle CR1: the first open reading frame of the turtle CR1 element encodes a protein with a novel zinc finger motif.
Mol. Biol. Evol. 14, 1206-17 (1997)
64. Okada N, and Hamada M.
The 3' ends of tRNA-derived SINEs originated from the 3' ends of LINEs: A new example from the bovine genome.
J. Mol. Evol. 44, Supplementally 52-6 (1997)
63. Ohshima K, Hamada M, Terai Y, and Okada N.
The 3' ends of tRNA-derived short interspersed repetitive elements are

derived from the 3' ends of long interspersed repetitive elements.
Mol. Cell. Biol. 16, 3756-64 (1996)

62. Kamaishi T, Hashimoto T, Nakamura Y, Nakamura F, Murata S, Okada N, Okamoto K, Shimizu M, and Hasegawa M.
Protein phylogeny of translation elongation factor EF-1a suggests microsporidians are extremely eukaryotes.
J. Mol. Evol. 42, 257-63 (1996)
61. Murata S, Takasaki N, Saitoh M, Tachida H, and Okada N.
Details of retropositional genome dynamics that provide a rationale for a generic division: The distinct branching of all the Pacific salmon and trout (*Oncorhynchus*) from the Atlantic salmon and trout (*Salmo*).
Genetics 142(3), 915-26 (1996)
60. Takasaki N, Park L, Kaeriyama M, Gharrett A J, and Okada N.
Characterization of species-specifically amplified SINEs in three salmonid species, chum salmon, pink salmon and kokanee: the local environment of the genome might be important for the generation of a dominant source gene at a newly retroposed locus.
J. Mol. Evol. 42(2), 103-16 (1996)
59. Kido Y, Saito M, Murata S, and Okada N.
Evolution of the source sequence of Hpa I short interspersed elements.
J. Mol. Evol. 41(6), 986-95 (1995)
58. Okada N, and Ohshima K.
Evolution of tRNA-derived SINEs. The Impact of short interspersed elements (SINEs) on the host genome.
Ed. Richard J. Maraia, Springer RG. Landes Company. (1995)
57. Matsuo M, Yokogawa T, Nishikawa K, Watanabe K, and Okada N.
Highly specific and efficient cleavage of squid tRNA(Lys) catalyzed by magnesium ions.
J. Biol. Chem. 270, 10097-104 (1995)
56. Matsuo M, Abe Y, Saruta Y, and Okada N.
Mollusk genes encoding lysine tRNA(UUU) contain introns.
GENE 165, 249-53 (1995)
55. Nameki N, Asahara H, Shimizu M, Okada N, and Himeno H.
Identity elements of *Saccharomyces cerevisiae* tRNA^{His}.
Nucleic Acids Res. 23, 389-94 (1995)

54. Nakanishi S, Ueda T, Hori H, Yamazaki N, Okada N, and Watanabe K.
A UGU sequence in the anticodon loop is a minimum requirement for recognition by *Escherichia coli* tRNA-guanine transglycosylase.
J. Biol. Chem. 269(51), 32221-5 (1994)
53. Ohshima K, and Okada N.
Generality of the tRNA origin of short interspersed repetitive elements (SINEs); Characterization of three different tRNA-derived retroposons in the octopus.
J. Mol. Biol. 243, 25-37 (1994).
52. Yokobori S, Hasegawa M, Ueda T, Okada N, Nishikawa K, and Watanabe K.
Relationship among Coelacanths, Lungfishes, and Tetrapods: A phylogenetic analysis based on mitochondrial cytochrome oxidase I gene sequences.
J. Mol. Evol. 38(6), 602-9 (1994)
51. Takasaki N, Murata S, Saitoh M, Kobayashi T, Park L, and Okada N.
Species-specific amplification of tRNA-derived SINEs via retroposition; a process of parasitization of entire genomes during the evolution of salmonids.
Proc. Natl. Acad. Sci. USA. 91(21), 10153-7 (1994) [OPEN ACCESS]
50. Kido Y, Himberg M, Takasaki N, and Okada N.
Amplification of distinct subfamilies of short interspersed elements (SINEs) during evolution of the Salmonidae.
J. Mol. Biol. 241(5), 633-44 (1994)
49. Sakagami M, Ohshima K, Mukoyama H, Yasue H, and Okada N.
A novel tRNA species as an origin of short interspersed repetitive elements (SINEs); Equine SINEs may have originated from tRNA(Ser).
J. Mol. Biol. 239(5), 731-5 (1994)
48. Yoshioka Y, Takahashi Y, Matsumoto S, Kojima S, Matsuoka K, Nakamura K, Ohshima K, Okada N, and Machida Y.
Mechanisms of T-DNA transfer and integration into plant chromosomes: role of virB, virD4 and virE2 and a short interspersed repetitive element (SINE) from tobacco.
Molecular Mechanisms of Bacterial Virulence, Kluwer Academic Publishers.
(eds. C. Kado & J. Crosa) 231-48 (1994)
47. Ozawa Y, Uchida K, Uchida M, Ami Y, Kushida S, Okada N, and Miwa M.

Isolation of cDNAs encoding the catalytic domain of poly(ADP-ribose) polymerase from *Xenopus laevis* and cherry salmon using heterologous oligonucleotide consensus sequences.

Biochem. Biophys. Res. Commun. 193, 119-25 (1993)

46. Murata S, Takasaki N, Saitoh M, and Okada N.
Determination of the phylogenetic relationships among Pacific salmonids using short interspersed elements (SINEs) as temporal landmarks of evolution.
Proc. Natl. Acad. Sci. USA .90(15), 6995-9 (1993) [OPEN ACCESS]
45. Yoshioka Y, Matsumoto S, Kojima S, Ohshima K, Okada N, and Machida Y.
Molecular characterization of a short interspersed repetitive element from tobacco that exhibits sequence homology to specific tRNAs.
Proc. Natl. Acad. Sci. USA. 90(14), 6562-6 (1993) [OPEN ACCESS]
44. Ohshima K, Koishi R, Matsuo M, and Okada N.
Several short interspersed repetitive elements (SINEs) in distant species may have originated from a common ancestral retrovirus: characterization of a squid SINE and a possible mechanism for generation of tRNA-derived retroposons.
Proc. Natl. Acad. Sci. USA .90(13), 6260-4 (1993) [OPEN ACCESS]
43. Kojima S, Hashimoto T, Hasegawa M, Murata S, Ohta S, Seki F, and Okada N.
Close phylogenetic relationship between Vestimentifera (tube worms) and Annelida revealed by the amino acid sequence of elongation factor-1a.
J. Mol. Evol. 37(1), 66-70 (1993)
42. Okada N, and Ohshima K.
A model for the mechanism of initial generation of short interspersed elements (SINEs).
J. Mol. Evol. 37(2), 167-70 (1993)
41. Kido Y, Aono M, Matsumoto K, Yamaki T, Saneyoshi M, and Okada N.
Shaping and reshaping of salmonid genomes by amplification of tRNA-derived retroposons during evolution.
Proc. Natl. Acad. Sci. USA.88(6), 2326-30 (1991) [OPEN ACCESS]
40. Okada N.
Transfer RNA-derived retroposons.
New Aspect of the Genetics of Molecular Evolution

(eds. M.Kimura and N.Takahata)
Japan Scientific Societies Press pp.227-41 (1991)

39. Okada N, Aono M, Endoh H, Kido Y, Koishi R, Matsumoto K, Matsuo M, Murata S, Nagahashi S, and Yamaki T.
Evolution of repetitive sequences. *Evolution of Life*.
(eds. S.Osawa and T.Honjo)
JSpringer-Verlag pp.175-86(1991)
38. Okada N.
SINEs: Short interspersed repeated elements of the eukaryotic genome.
Trends in Ecology & Evolution. 6(11), 358-61 (1991)
37. Okada N.
SINEs.
Current Opinion in Genetics & Development 1, pp.498-504 (1991)
36. Koishi R, and Okada N.
Distribution of the salmonid HPA-1 family in the salmonid species demonstrated by In vitro run-off transcription assay of total genomic DNA : A procedure to estimate repetitive frequency and sequence divergence of a certain repetitive family with a few known sequences.
J. Mol. Evol. 32(1), 43-52 (1991)
35. Nagahashi S, Endoh H, Suzuki Y, and Okada N.
Characterization of a tandemly repeated DNA sequence family originally derived by retroposition of tRNAGlu in the newt.
J. Mol. Biol. 222, 391-404 (1991)
34. Endoh H, Nagahashi S, and Okada N.
Highly repetitive and transcribable sequence in the tortoise genome is probably a retroposon.
Eur. J. Biochem. 189(1), 25-31 (1990)
33. Okada N.
Transfer RNA-like structure of the human Alu family: Implications of its generation mechanism and possible functions.
J. Mol. Evol. 31(6), 500-10 (1990)
32. Matsumoto K, Takii T, and Okada N.
Characterization of a new termination signal for RNA polymerase III responsible for generation of a discrete-sized RNA transcribed from salmon genomic total DNA in a HeLa cell extract.
J. Biol. Chem. 264(2),1124-31 (1989)

31. Endoh H, Nagahashi S, and Okada N.
Tetrahymena pyriformis DNA fragment with a gene cluster for 3 putative serine tRNAs and an asparagine tRNA.
Nucleic Acids Res. 17, 10122 (1989)
30. Okada N, Aono M, Matsumoto K, Kido Y, Koishi R, and Saneyoshi M.
Shaping and reshaping of the genomes of salmonid species during evolution. Biology of Charrs and Masu Salmon .
(eds. Kawanabe, Yamazaki and Noakes)
Special Vol. 1, Physiology and Ecology Japan, pp.451-4 (1989)
29. Okada N, Aono M, Matsumoto K, Endoh H, Koishi R, Takii T, and Suzuki Y.
Several aspects of total genomic DNA transcription in a HeLa cell extract.
Nucleic Acids Res. Sympo. Ser. No. 19, 181-4 (1988)
28. Hirono M, Endoh H, Okada N, Numata O, and Watanabe Y.
Tetrahymena actin, cloning and sequencing of the Tetrahymena actin gene and identification of its gene product.
J. Mol. Biol. 194, 181-92 (1987)
27. Okada N, Mimori T, Mukai R, Kashiwagi H, and Hardin J A.
Characterization of human autoantibodies that selectively precipitate the 7SL RNA component of the signal recognition particle.
J. Immunology 138, 3219-23 (1987)
26. Endoh H , and Okada N.
Total DNA transcription in vitro: A procedure to detect highly repetitive and transcribable sequences with tRNA-like structures.
Proc. Natl. Acad. Sci. USA 83, 251-5 (1986)
25. Matsumoto K, Murakami K, and Okada N.
Gene for lysine tRNA^{L1} as a progenitor of the highly repetitive and transcribable sequences present in the salmon genome.
Proc. Natl. Acad. Sci. USA 83, 3156-60 (1986)
24. Sakamoto K, and Okada N.
Rodent type 2 Alu family, rat identifier sequence, rabbit C family, and bovine or goat 73-bp repeat may have evolved from tRNA genes.
J. Mol. Evol. 22, 134-40 (1985)
23. Okada N, Endoh H, Sakamoto K, and Matsumoto K.
Many highly repetitive and transcribable sequences are derived from

tRNA genes.

Proc. Jpn. Acad. 61, Ser. B, 363-7 (1985)

22. Sakamoto K, and Okada N.
5-Methylcytidylic modification of in vitro transcript from the rat identifier sequence; evidence that the transcript forms a tRNA-like structure.
Nucleic Acids Res. 13, 7195-206 (1985)
21. Hoshino H, Tanaka H, Shimotono K, Miwa M, Okada N, and Sugimura T.
Nucleotide sequences homologous to human T-cell leukemia virus are present in pig and wild bore cells.
Gann. 75, 849-52 (1984)
20. Okada N, Mukai R, Harada F, Kabashima T, Nakao Y, Yamane K, Ohshima Y, Sakamoto K, Itoh M, Kashiwagi H, and Hamaguchi H.
Isolation of a novel antibody which precipitates ribonucleoprotein complex containing threonine tRNA from a patient with polymyositis.
Eur. J. Biochem. 139, 425-9 (1984)
19. Sakamoto K, Komimani R, Mishima Y, and Okada N.
The 6S RNA transcribed from rodent total DNA in vitro is the transcript of the type 2 Alu family.
Mol. Gen. Genet. 194, 1-6 (1984)
18. Matsumoto K, Murakami K, and Okada N.
Pseudouridylic modification of a 6S RNA transcribed in vitro from highly repetitive and transcribable (Hirt) sequences of salmon total DNA.
Biochem. Biophys. Res. Commun. 124, 514-22 (1984)
17. Okada N, Sakamoto K. and Kondo A.
Total DNA transcription reveals the existence of highly repetitive transcribable sequences in higher animals.
J. Biochem. 93, 723-31 (1983)
16. Watanabe-Nagasu N., Itoh Y., Tani T., Okano K., Koga N., Okada N. and Ohshima Y.
Structural analysis of gene loci for rat U1 small nuclear RNA.
Nucleic Acids Research. 11, 1791-801 (1983)
15. Tani T., Watanabe-Nagasu N., Okada N. and Ohshima Y.
Molecular cloning and characterization of a gene for rat U2 small nuclear RNA.
J. Mol. Biol. 168, 579-94 (1983)

14. Okada N., Sakamoto K., Itoh Y. and Ohshima Y.
Sequence determination of rat U5 RNA using a chemical modification procedure for counteracting sequence compression.
J. Biochem. 91, 1281-91 (1982)
13. Nakao Y., Mukai R., Kabashima T., Ohshima Y., Hamaguchi H., Kashiwagi H. and Okada N.
A novel antibody which precipitates 7.5S RNA is isolated from a patient with autoimmune disease.
Biochem. Biophys. Res. Commun. 109, 1332-8 (1982)
12. Shindo-Okada N., Kuchino Y., Harada F., Okada N. and Nishimura S.
Biological and structural differences between tRNA^{Val} species isolated from rat ascites hepatoma cells and normal rat liver.
J. Biochem. 90, 535-44 (1981)
11. Ohshima Y., Itoh M., Okada N. and Miyata T.
Novel models for RNA splicing that involve a small nuclear RNA.
Proc. Natl. Acad. Sci. USA. 78, 4471-4 (1981)
10. Ohshima Y., Okada N., Tani T., Itoh Y. and Itoh M.
Nucleotide sequences of mouse genomic loci including a gene or pseudogene for U6(4.8S) nuclear RNA.
Nucleic Acids Res. 9, 5145-58 (1981)
9. Shindo-Okada N., Okada N., Ohgi T., Goto T. and Nishimura S.
Transfer ribonucleic acid guanine transglycosylase isolated from rat liver.
Biochemistry 19, 395-400 (1980)
8. Okada N. and Nishimura S.
Isolation and characterization of a guanine insertion enzyme, a specific tRNA transglycosylase, from *Escherichia coli*.
J. Biol. Chem. 254, 3061-6 (1979)
7. Okada N., Noguchi S., Kasai H., Shindo-Okada N., Ohgi T., Goto T. and Nishimura S.
Novel mechanism of posttranscriptional modification of tRNA. - insertion of bases of Q-precursors into transfer-RNA by a specific transfer-RNA transglycosylase reaction.
J. Biol. Chem. 254, 3067-73 (1979)
6. Okada N., Noguchi S., Nishimura S., Ohgi T., Goto T., Crain P.F. and MacCloskey J.A.
Structure determination of a nucleoside Q precursor isolated from *E. coli*

tRNA: 7-(aminomethyl)-7-deazaguanosine.
Nucleic Acids Res. 5, 2289-96 (1978)

5. Okada N., Shindo-Okada N., Sato S., Itoh Y.H., Oda K-I. and Nishimura S.
Detection of unique tRNA species in tumor tissues by Escherichia coli guanine insertion enzyme.
Proc. Natl. Acad. Sci. USA 75, 4247-51 (1978)
4. Okada N. Shindo-Okada N. and Nishimura S.
Isolation of mammalian tRNA^{Asp} and tRNA^{Tyr} by lectin-Sepharose affinity column chromatography.
Nucleic Acids Res. 4, 415-23 (1977)
3. Okada N. and Nishimura S.
Enzymatic synthesis of Q* nucleoside containing mannose in the anticodon of tRNA: Isolation of a novel mannosyltransferase from a cell-free extract of rat liver.
Nucleic Acids Res. 4, 2931-7 (1977)
2. Okada N., Yasuda T. and Nishimura S.
Detection of nucleoside Q precursor in methyl-deficient E. coli tRNA.
Nucleic Acids Res. 4, 4063-75 (1977)
1. Okada N., Harada F. and Nishimura S.
Specific replacement of Q base in the anticodon of tRNA by guanine catalyzed by a cell-free extract of rabbit reticulocytes.
Nucleic Acids Res. 3, 2593-603 (1976)